

Gualaquiza and Cuenea. "Tortola." "Irides hazel, cere bluish flesh-colour; rim round the eyes greenish yellow; bill black; legs red."

82. **COLUMBULA CRUZIANA**, D'Orb.; Bp. Conspl. ii. p. 80.

Cuenea, 1 ex. ♀. Beak black; cere yellow; irides red; legs red, claws black: contents of stomach very small seeds. 'Tortolita,' Spanish. "I saw nine together on the tiles of a house in Cuenea; also I observed it in the fields, sometimes in threes and fours, often solitary."

83. **GLOTTIS MELANOLEUCA** (Gm.).

Cuenea. Eyes black; bill green; legs deep orange.

84. **TRINGOIDES MACULARIUS** (Linn.).

Gualaquiza, ♀, 2 ex. Young birds, but the spots below are just discernible in one specimen.

85. **TOTANUS SOLITARIUS**, Wilson.

Gualaquiza, 1 ex. ♀. Irides dark hazel; bill blackish; legs dull green. Frequents the stones and margins of the river.

86. **TRINGA** — ?

Cuenea, Oct. 1857.

87. **TIGRISOMA TIGRINUM**.

Gualaquiza, Feb. 1858.

3. **SYNOPSIS GENERIS FRINGILLINI ERYTHRURÆ, AUCTORE GUSTAVO HARTLAUB, MED. DOCT.**

a. **SPECIES TYPICÆ.**

1. **ERYTHRURA PRASINA** (Sparm.). *Viridis, fronte guttureque dilute cyaneis, abdomine medio et caudu rubris.*

Hab. In Java et Sumatra.

Syn. *Loxia prasina*, Sparm. Mus. Carls. t. 72, 73.—*Emberiza quadricolor*, Gm. L. no. 65.—*Emberiza cyanopis*, Gm. ibid. no. 66.—*Fringilla sphenura*, Temm. Pl. Col. 96.—*Erythrura viridis*, Sw. Confer Bp. Conspl. p. 457.

2. **ERYTHRURA TRICHOA** (Kittl.). *Læte viridis; facie et regione parotica pulchre cyaneis; canda breviuscula cum tectricibus superioribus rubra, rectricibus intermediis cernulescentibus.*

Hab. In ins. Ualan Carolinarum.

Syn. *Fringilla trichrou*, v. Kittl. Mem. Acad. Pet. 1835, ii. p. 8. t. 10, fig. bona.—*Erythrura trichroa*, Bp. Conspl. i. p. 457; v. Kittl. Denkwürdigk. ii. p. 38.

3. **ERYTHRURA TRICOLOR** (Vieill.). *Læte viridis; fronte, genis et gastræo toto læte cyaneis; cauda parum elongata rubra.*
Hab. Timor (Mus. Paris. et Lugd.).
Syn. *Fringilla tricolor*, Vieill. Ois. Chant. pl. 20; id. Encyclop. p. 974.—*Erythrura tricolor*, Bp. Conspl. p. 457.
4. **ERYTHRURA PSITTACEA** (Gm.). *Læte viridis; fronte, sincipite, gutture, uropygio et cauda coccineo-sanguineis.*
Hab. In Nova Caledonia.
Syn. *Parrot Finch*, Lath. Gen. Syn. iii. 287. pl. 48; id. G. H. vi. p. 81. pl. 96.—*Fringilla psittacea*, Gm. S. N. i. 903.—*Char donneret acalanthe*, Vieill. Ois. Chant. pl. 32; id. Encycl. p. 982.—*Fringilla pulchella*, R. Forster, Deser. Anim., ed. Licht. p. 273.—*Erythrura psittacea*, Bp. Conspl. p. 457.—*Poëphila Paddoni*, Mac gillivr. Ann. & Mag. N. H. 1858, p. 273.
5. **ERYTHRURA PUCHERANII**, Bp. *Glauco-cyanea; pileo genis tectricibusque caudæ superioribus sanguineis.*
Hab. In insulis Oceaniæ.
Syn. *Erythrura Pucheranii*, Bp. Conspl. i. p. 457.
6. **ERYTHRURA CYANOVIRENS** (Peale). *Viridis, nitore glaucescente: capite coccineo: gutture pectoreque dilute cyaneis: tectricibus duabus intermediis totis rubentibus, reliquis in medio fuscescenscentibus in margine rubentibus.*
Hab. In. ins. Navigatorum Upolu.
Syn. *Geospiza cyanovirens*, Peale, Unit. St. Expl. Exp. Ornith. p. 117; Hartl. in Wieg. Archiv. xviii. p. 104.
7. **ERYTHRURA PEALII**, nob. *Læte viridis; pileo, regione parotica, et tectricibus caudæ superioribus scarlatinis; pectore pulchre cyaneo, gulam nigricantem versus saturate indigotico; rectricibus nigricantibus viridi marginatis.*
Hab. In insulis Fee-jee.
Syn. *Geospiza prasina*, Peale, Unit. St. Expl. Exp. Orn. p. 116.—*Erythrura pealii*, Hartl. Wieg. Arch. xviii. p. 104.
- SP. MINUS TYPICA.
8. **ERYTHRURA PHAETON** (Hombr. & Jacq.), Puch. Voy. au Pol. Sud. Ois. p. 99.—*Neochmia phaeton*, Hombr. & Jacq.; Bp. Conspl. i. p. 458; Gould, B. Austr. iii. pl. 83; Reichenb. Vög. Neuholl. p. 115.
Hab. In Australia Boreali.
4. NOTES ON THE HABITS OF THE SCYTHROPS NOVÆ HOLLANDIÆ. BY GEORGE BENNETT, ESQ., CORR. MEMB. ZOOL. SOC. COMMUNICATED TO JOHN GOULD, ESQ., F.R.S., V.P.Z.S., ETC., IN A LETTER DATED SYDNEY, JUNE 8, 1858.
- “ I send you a few notes on the *Scythrops Novæ Hollandiæ*, or